

THE OPTICAL GRAVITATIONAL LENSING EXPERIMENT. A HUNT FOR CAUSTIC CROSSINGS IN QSO 2237+0305.*

P.R. WOŹNIAK¹, A. UDALSKI², M. SZYMAŃSKI², M. KUBIAK², G. PIETRZYŃSKI², I. SOSZYŃSKI²,
 AND K. ŻEBRUŃ².

¹Princeton University Observatory, Princeton, NJ 08544-1001, USA
 e-mail: wozniak@astro.princeton.edu

²Warsaw University Observatory, Al. Ujazdowskie 4, 00-478 Warszawa, Poland
 e-mail: udalski,msz,mk,pietrzyn,soszynsk,zebrun@sirius.astrouw.edu.pl

Draft version February 1, 2008

ABSTRACT

In 1998 The Optical Gravitational Lensing Experiment (OGLE) successfully implemented automated data reductions for QSO 2237+0305. Using a new image subtraction method we achieved a differential photometry scatter of 1–5% for images A–D respectively. Combined with a time sampling of 1–2 times a week this is sufficient for early detection of caustic crossings. Nearly real time photometry of QSO 2237+0305 is available from the OGLE web site <http://www.astro.princeton.edu/~ogle/ogle2/huchra.html>. During the 1999 observing season, the apparent *V* magnitude of the A, B, C and D images changed by 0.50, 0.15, 0.65 and 0.35 mag, respectively. Most likely however, none of the microlensing events involved a caustic crossing. The most rapid variation was 0.25 mag in 30 days, observed for image C. The alert system will continue to be active in the next observing season from late April until September 2000, when OGLE suspends operation for an upgrade. Observations will resume for season of 2001.

Subject headings: galaxies: photometry – gravitational lensing – quasars: individual (Q2237+0305)

1. INTRODUCTION

The Optical Gravitational Lensing Experiment is a long term project focusing on detection and monitoring of microlensing events in crowded fields. During the second phase of the experiment, data are collected with the dedicated 1.3 m Warsaw Telescope at the Las Campanas Observatory (LCO), Chile (Udalski, Kubiak and Szymański 1997). In 1997, OGLE started a monitoring program for two multiply imaged lensed quasars: QSO 2237+0305 and HE 1104-1805. In QSO 2237+0305 (Huchra’s lens) one expects individual quasar images to vary due to microlensing in the bulge of the macrolensing galaxy (Webster et al. 1991, Schneider et al. 1988). Theoretical models predict high magnification events at caustic crossings. The sharpest features in the light curve provide in principle a size measurement of the source, in this case a quasar (e.g. Wambsganss et al. 1990). This letter presents observations made in 1999, previously unpublished except for a web site release. Also we would like to call the attention of astronomical community to the fact that good photometry of QSO 2237+0305 is available practically in real time, and encourage follow up programs.

2. OBSERVATIONS

The OGLE monitoring of Huchra’s lens is being carried out with the 1.3 m Warsaw Telescope at the Las Campanas Observatory, Chile, which is operated by the

Carnegie Institution of Washington. The “first generation” camera uses a SITe 2048 × 2048 CCD detector with 24 μ m pixels resulting in 0.417 arcsec/pixel scale. Images of QSO 2237+0305 are taken in normal mode (still frame) at “medium” readout speed with the gain 7.1 e^- /ADU and readout noise of 6.3 e^- . For the details of the instrumentation setup, we refer the reader to Udalski, Kubiak and Szymański (1997).

Good seeing is essential for ground based measurements of Huchra’s lens. Therefore, observations are restricted to nights with seeing better than about 1.4 arcsec, preferably with low sky background. The median FWHM of the seeing disk was 1.3 arcsec in the data presented here. In satisfactory weather conditions between late April and mid December, when QSO 2237+0305 is visible from LCO, two 4 minute exposures in the standard *V* photometric band are taken, shifted by few arc seconds with respect to each other. This can be done typically once or twice per week without a significant slow down of the primary program.

3. ONLINE REDUCTIONS

The method which enabled reliable difference image photometry of QSO 2237+0305 was developed by Alard and Lupton (1998) and further generalized for spatially variable kernels by Alard (1999). By taking the difference of each frame with the reference image (a stack of 20 best frames) one avoids many complications associated with the

*Based on observations obtained with the 1.3 m Warsaw Telescope at the Las Campanas Observatory of the Carnegie Institution of Washington.




FIG. 1.— Light curve of QSO 2237+0305. Also shown is the light curve of the 18.14 mag reference star, shifted by 1.1 mag for clarity.

presence of a barred spiral and the small separations of quasar images. The photometric pipeline consists of a set of stand-by programs for detections of stars, registration of frames with the reference frame, actual subtraction and fitting a 4 PSF model of Huchra's lens. A detailed description of the implementation can be found in Woźniak et al. (2000). The OGLE II data pipeline automatically detects new frames and performs bias and flat-field corrections (Udalski, Kubiak, & Szymański 1997). Initially reduced frames are passed to image subtraction programs. Within 10 minutes after exposure a new photometric point is e-mailed and can be posted on the OGLE web page. However for practical reasons new data usually appear on the web the next morning.

4. LIGHT CURVE

Photometry of the four components of QSO 2237+0305 is shown in Fig. 1. The light curve of the 18.14 mag comparison star is also included. The star was measured with exactly the same method as quasar images. Wyithe et al. (2000) interpret these observations. Based on their models we may expect a caustic crossing in the near future. In the upcoming 2000 observing season for QSO 2237+0305, the time sampling of our photometry should be comparable to the season of 1999. After September 2000 OGLE will suspend operation for an upgrade. Observations will resume in season of 2001. Machine readable data can be obtained from the OGLE web page at <http://www.astro.uw.edu.pl/~ftp/ogle> and its USA mirror <http://www.astro.princeton.edu/~ogle>.

This work was supported with NSF grant AST-9820314 to Bohdan Paczyński.

REFERENCES

- Alard, C., & Lupton, R. H., 1998, *ApJ*, 503, 325
- Alard, C., 1999, *A&A*, submitted (= astro-ph/9903111)
- Schneider, D. P., Turner, E. L., Gunn, J. E., Hewitt, J. N., Schmidt, M., & Lawrence, C. R., 1988, *AJ*, 95, 1619
- Udalski, A., Kubiak, M., & Szymański, M., 1997, *Acta Astron.*, 47, 319
- Wambsganss, J., Paczyński, B., & Schneider, P., 1990, *ApJ*, 358, L33
- Webster, R. L., Ferguson, A. M. N., Corrigan, R. T., & Irwin, M. J., 1991, *AJ*, 102, 1939
- Woźniak, P. R., Alard, C., Udalski, A., Szymański, M., Kubiak, M., Pietrzyński, G., Żebruń, K., 2000, *ApJ*, 529, 88
- Wyithe, J. S. B., Turner, E. L., Webster, R. L., 2000, *MNRAS*, submitted (astro-ph/0001307)